



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

Enzymatic Assay of CHONDROITINASE ABC (EC 4.2.2.4)

PRINCIPLE:



CONDITIONS: T = 37°C, pH = 8.0, A232nm, Light path = 1 cm

METHOD: Spectrophotometric Stop Rate Determination

REAGENTS:

- A. 250 mM Tris HCl and 300 mM Sodium Acetate Buffer with 0.05% (w/v) Bovine Serum Albumin, pH 8.0 at 37°C
(Prepare 200 ml in deionized water using Trizma Base, Sodium Acetate, Trihydrate, and Albumin, Bovine. Adjust to pH 8.0 at 37°C with 1 M HCl or 1 M NaOH.)
- B. 0.5% (w/v) Chondroitin Sulfate A Solution (Chon A)
(Prepare 5 ml in Reagent A using Chondroitin Sulfate A, Sodium Salt.)
- C. 0.5% (w/v) Chondroitin Sulfate B Solution (Chon B)
(Prepare 5 ml in Reagent A using Chondroitin Sulfate B, Sodium Salt.)
- D. 0.5% (w/v) Chondroitin Sulfate C Solution (Chon C)
(Prepare 5 ml in Reagent A using Chondroitin Sulfate, Sodium Salt, from shark cartilage.)
- E. 0.01% (w/v) Bovine Serum Albumin Solution (BSA)
(Prepare 20 ml in deionized water using Albulrin Bovine Serum.)
- F. 50 mM Potassium Chloride Solution, pH 1.8 at 25°C (KCl)
(Prepare 50 ml in deionized water using Potassium Chloride. Adjust to pH 1.8 at 25°C with 1 M HCl.)



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

REAGENTS: (continued)

- G. Chondroitinase ABC Enzyme Solution
(Immediately before use, prepare a solution containing 0.06 - 0.10 unit/ml of Chondroitinase ABC in cold Reagent E.)

PROCEDURE:

Pipette (in milliliters) the following reagents into suitable tubes:

	<u>Test 1</u>	<u>Test 2</u>	<u>Test 3</u>
Reagent G (Enzy Soln)	0.80	0.80	0.80
Equilibrate to 37°C for 2 minutes. Then add:			
Reagent B (Chon A)	0.20	—	—
Reagent C (Chon B)	—	0.20	—
Reagent D (Chon C)	—	—	0.20

Mix by inversion. Incubate at 37°C for 21 minutes. At 0, 3, 6, 9, 12, 15, 18, and 21 minutes (T_T), transfer 0.10 ml from each Test to separate tubes containing 0.90 ml of Reagent F (KCl). The Blank for this assay is T_0 .

Incubate each tube for an additional 10 minutes at 37°C. Centrifuge for 10 minutes and transfer the Test and Blank supernatants to suitable quartz cuvettes. Record the absorbance at 232 nm for each tube.

CALCULATIONS:

T_T = Time intervals (minutes) for enzyme assay incubation

$$\Delta A_{232\text{nm}} \text{ Test } (T_T) = A_{232\text{nm}} \text{ Test } (T_T) - A_{232\text{nm}} \text{ Blank } (T_0)$$

Plot the $\Delta A_{232\text{nm}}$ Test (T_T) vs. Time. Determine the $\Delta A_{232\text{nm}}/\text{min}$ Test from a linear portion of the graph.



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

CALCULATIONS:

$$\text{Units/ml} = \frac{(\Delta A_{232\text{nm}}/\text{min Test}) (1.0) (\text{df})}{(\text{E}_{\text{mM}}) (0.10) (0.8)}$$

df = Dilution factor

E_{mM} = Millimolar extinction coefficient of unsaturated disaccharides: 5.1 for products from Chondroitin A and Chondroitin B, and 5.5 for products from Chondroitin C.

0.10 = Volume (in milliliter) of reaction mix used

T = Time (in minutes) of the assay as per the Unit Definition

1.0 = Total volume (in milliliter) of assay

0.8 = Volume (in milliliter) of enzyme used

$$\text{Unit/mg solid} = \frac{\text{units/ml enzyme}}{\text{mg solid/ml enzyme}}$$

UNIT DEFINITION:

One unit will liberate 1.0 μ mole of 2-acetamido-2-deoxy-3-O- (β -D-gluc-4-ene-pyranosyluronic acid)-4-O-sulfo-D-galactose from chondroitin sulfate A or 1.0 μ mole of 2-acetamido-2-deoxy-3-O-(β -D-gluc-4-ene-pyranosyluronic acid)-6-O-sulfo-D-galactose from chondroitin sulfate C per minute at pH 8.0 at 37°C.

FINAL ASSAY CONCENTRATION:

In a 1.00 ml reaction mix, the final concentrations are 50 mM Tris, 60 mM sodium acetate, 0.02% (w/v) bovine serum albumin, 0.1% (w/v) chondroitin sulfate A, B or C, and 0.048 - 0.08 unit chondroitinase ABC.

REFERENCE:

Saito, H., Yamagata, T. and Suzuki, S. (1968) *J. Biol. Chem.* **243**, 1536-1542

Yamagata, T., Saito, H., Habuchi, O., and Suzuki, S. (1968) *J. Biol. Chem.* **243**, 1523-1535

NOTES:

1. This assay is based on the cited references.