

产品名称: **VE-821**

产品别名: **VE-821**

生物活性:					
<b>Description</b>	VE-821 is a potent ATP-competitive inhibitor of ATR with $K_i/IC_{50}$ of 13 nM/26 nM.				
<b>IC<sub>50</sub> &amp; Target</b>	ATR	ATM	DNA-PK	PI3K $\gamma$	
	13 nM (K <sub>i</sub> )	16 $\mu$ M (K <sub>i</sub> )	2.2 $\mu$ M (K <sub>i</sub> )	3.9 $\mu$ M (K <sub>i</sub> )	
<b>In Vitro</b>	VE-821 shows excellent selectivity for ATR with minimal cross-reactivity against the related PIKKs ATM, DNA-PK, mTOR and PI3K $\gamma$ (K <sub>i</sub> s of 16 $\mu$ M, 2.2 $\mu$ M, >1 $\mu$ M and 3.9 $\mu$ M, respectively) and against a large panel of unrelated protein kinases[1]. VE-821 (compound 27) also inhibits ATM and DNA-PK with IC <sub>50</sub> of >8 $\mu$ M, and 4.4 $\mu$ M, respectively[2]. VE-821 significantly enhances the sensitivity of PSN-1, MiaPaCa-2 and primary PancM pancreatic cancer cells to radiation and Gemcitabine under both normoxic and hypoxic conditions. ATR inhibition by VE-821 leads to inhibition of radiation-induced G <sub>2</sub> /M arrest in cancer cells. In both PSN-1 and MiaPaCa-2 cells, 1 $\mu$ M VE-821 inhibits phosphorylation of Chk1 (Ser 345) after treatment with Gemcitabine (100 nM), radiation (6 Gy) or both, at 2 h post-irradiation[3].				
<b>Solvent&amp;Solubility</b>	<b>In Vitro:</b> DMSO : 50 mg/mL (135.72 mM; Need ultrasonic) H <sub>2</sub> O : < 0.1 mg/mL (insoluble)				
	<b>Preparing Stock Solutions</b>	<b>Solvent</b> \ <b>Mass</b> \ <b>Concentration</b>	<b>1 mg</b>	<b>5 mg</b>	<b>10 mg</b>
		1 mM	2.7144 mL	13.5718 mL	27.1437 mL
		5 mM	0.5429 mL	2.7144 mL	5.4287 mL
	10 mM	0.2714 mL	1.3572 mL	2.7144 mL	
*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液; 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80°C, 6 months; -20°C, 1 month。-80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。					
<b>In Vivo:</b> 1.VE-821 is prepared in vehicle (10% PEG300, 2.5% Tween-80, pH 4)[4].					
<b>References</b>	[1]. Reaper PM, et al. Selective killing of ATM- or p53-deficient cancer cells through inhibition of ATR. <i>Nat Chem Biol.</i> 2011 Apr 13;7(7):428-30. [2]. Charrier JD, et al. Discovery of potent and selective inhibitors of ataxia telangiectasia mutated and Rad3 related (ATR) protein kinase as potential anticancer agents. <i>J Med Chem.</i> 2011 Apr 14;54(7):2320-30. [3]. Prevo R, et al. The novel ATR inhibitor VE-821 increases sensitivity of pancreatic cancer cells to radiation and chemotherapy. <i>Cancer Biol Ther.</i> 2012 Sep;13(11):1072-81. [4]. Muralidharan SV, et al. BET bromodomain inhibitors synergize with ATR inhibitors to induce DNA damage, apoptosis, senescence-associated secretory pathway and ER stress in Myc-induced lymphoma cells. <i>Oncogene.</i> 2016 Sep 8;35(36):4689-97.				
实验参考:					
<b>Cell Assay</b>	MiaPaCa-2, PSN-1 and Panc1 cells (5×10 <sup>4</sup> ) are plated in 96-well plates and after 4 h treated with increasing concentrations of VE-821 at 1 h before irradiation with a single dose of 4 Gy. Medium is replaced 72 h post-irradiation at which point viability is measured using the using the Alamar Blue				

	<p>assay. Cells are allowed to proliferate and cell viability is again analyzed at day 10 for the different treatment conditions. Cell viability and surviving fraction are normalized to the untreated (control) group[3].</p>
<b>Kinase Assay</b>	<p>The ability of compounds (e.g., VE-821) to inhibit ATR, ATM or DNAPK kinase activity is tested using a radiometric-phosphate incorporation assay. A stock solution is prepared consisting of the appropriate buffer, kinase, and target peptide. To this is added the compound of interest, at varying concentrations in DMSO to a final DMSO concentration of 7%. Assays are initiated by addition of an appropriate [<math>^3\text{H}</math>]ATP solution and incubated at 25°C. Assays are stopped, after the desired time course, by addition of phosphoric acid and ATP to a final concentration of 100 mM and 0.66 <math>\mu\text{M}</math>, respectively. Peptides are captured on a phosphocellulose membrane, prepared, and washed six times with 200 <math>\mu\text{L}</math> of 100 mM phosphoric acid, prior to the addition of 100 <math>\mu\text{L}</math> of scintillation cocktail and scintillation counting on a 1450 Microbeta Liquid Scintillation Counter. Dose-response data are analyzed using GraphPad Prism software[2].</p>
<b>References</b>	<p>[1]. Reaper PM, et al. Selective killing of ATM- or p53-deficient cancer cells through inhibition of ATR. <i>Nat Chem Biol.</i> 2011 Apr 13;7(7):428-30.</p> <p>[2]. Charrier JD, et al. Discovery of potent and selective inhibitors of ataxia telangiectasia mutated and Rad3 related (ATR) protein kinase as potential anticancer agents. <i>J Med Chem.</i> 2011 Apr 14;54(7):2320-30.</p> <p>[3]. Prevo R, et al. The novel ATR inhibitor VE-821 increases sensitivity of pancreatic cancer cells to radiation and chemotherapy. <i>Cancer Biol Ther.</i> 2012 Sep;13(11):1072-81.</p> <p>[4]. Muralidharan SV, et al. BET bromodomain inhibitors synergize with ATR inhibitors to induce DNA damage, apoptosis, senescence-associated secretory pathway and ER stress in Myc-induced lymphoma cells. <i>Oncogene.</i> 2016 Sep 8;35(36):4689-97.</p>

源叶生物