



上海源叶生物科技有限公司
 Shanghai yuanye Bio-Technology Co., Ltd
 电话: 021-61312973 传真: 021-55068248
 网址: www.shyuanye.com
 邮箱: shyysw@sina.com

产品名称: **SAR245409**
 产品别名: **Voxtalisib**

| 生物活性: | | | | | | |
|-------------------------------------|--|----------------------------|---------------------------|----------------------------|----------------------------|----------------------------|
| Description | Voxtalisib (XL765) is a potent PI3K inhibitor, which has a similar activity toward class I PI3K (IC ₅₀ s=39, 113, 9 and 43 nM for p110α, p110β, p110γ and p110δ, respectively), also inhibits DNA-PK (IC ₅₀ =150 nM) and mTOR (IC ₅₀ =157 nM). Voxtalisib (XL765) inhibits mTORC1 and mTORC2 with IC ₅₀ s of 160 and 910 nM, respectively. | | | | | |
| IC₅₀ & Target | p110γ | p110α | p110δ | p110β | mTOR | mTORC1 |
| | 9 nM (IC ₅₀) | 39 nM (IC ₅₀) | 43 nM (IC ₅₀) | 113 nM (IC ₅₀) | 157 nM (IC ₅₀) | 160 nM (IC ₅₀) |
| | mTORC2 | DNA-PK | | | | |
| | 910 nM (IC ₅₀) | 150 nM (IC ₅₀) | | | | |
| In Vitro | Voxtalisib (XL765) displays potent inhibitory activity against class I PI3K isoforms p110α, p110β, p110δ, and p120γ, with IC ₅₀ s of 39, 110, 43, and 9 nM, respectively. The IC ₅₀ value for inhibition of PI3Kα by Voxtalisib is determined at various concentrations of ATP, revealing Voxtalisib be an ATP-competitive inhibitor with an equilibrium inhibition constant (K _i) value of 13 nM. Voxtalisib also inhibits mTOR (IC ₅₀ s of 160 and 910 nM for mTORC1 and mTORC2, respectively) in an immune-complex kinase assay and the PI3K-related kinase DNA-PK (IC ₅₀ value of 150 nM). In contrast, Voxtalisib (XL765) has relatively weak inhibitory activity toward the class III PI3K vacuolar sorting protein 34 (VPS34; IC ₅₀ value of ~9.1 μM). Consistent with its inhibitory activity against purified PI3K proteins, SAR245409 inhibits EGF-induced PIP3 production in PC-3 and MCF7 cells with IC ₅₀ s of 290 and 170 nM, respectively. The ability of Voxtalisib to inhibit phosphorylation of key signaling proteins downstream of PI3K is examined by assessing its effects on EGF-stimulated phosphorylation of AKT and on nonstimulated phosphorylation of S6 in PC-3 cells by cell-based ELISA. Voxtalisib inhibits these activities with IC ₅₀ s of 250 and 120 nM, respectively. In MCF7 and PC-3 cells, Voxtalisib inhibits proliferation (monitored by BrdUrd incorporation) with IC ₅₀ s of 1,070 and 1,840 nM, respectively. To further characterize the effects of Voxtalisib on tumor cell growth, an assay monitoring the anchorage-independent growth of PC-3 and MCF7 cells in soft agar over a 14-day period is used. SAR245409 inhibits colony growth with an IC ₅₀ value of 270 nM in PC-3 cells and 230 nM in MCF7 cells[2]. | | | | | |
| In Vivo | Oral administration of Voxtalisib (XL765) causes a dose-dependent decrease of phosphorylation of AKT, p70S6K, and S6 in the tumors, reaching a maximum of 84% inhibition of S6 phosphorylation at 30 mg/kg at 4 hours. The dose-response relationships derive from the 4 hours time point predict 50% inhibition of AKT, p70S6K, and S6 phosphorylation to occur at doses of 19 mg/kg (pAKT ^{T308} and pAKT ^{S473}), 51 mg/kg (p-p70S6K), and 18 mg/kg (pS6). Inhibition of AKT, p70S6K, and S6 phosphorylation in MCF7 tumors following a 30 mg/kg dose of Voxtalisib is maximal at 4 hours, reaching 61% to 84%; however, the level of inhibition decreases to 0% to 42% by 24 hours, and minimal or no inhibition is evident by 48 hours. Following a 100 mg/kg dose of Voxtalisib, inhibition is also maximal at 4 hours (52%-75%)[2]. | | | | | |
| | In Vitro: DMSO : 16 mg/mL (59.20 mM; Need ultrasonic and warming) | | | | | |
| | | Solvent Concentration | Mass | 1 mg | 5 mg | 10 mg |



上海源叶生物科技有限公司
 Shanghai yuanye Bio-Technology Co., Ltd
 电话: 021-61312973 传真: 021-55068248
 网址: www.shyuanye.com
 邮箱: shyysw@sina.com

| | | | | | |
|--|---|-----------|-----------|------------|------------|
| Solvent&Solubility | Preparing Stock Solutions | | | | |
| | | 1 mM | 3.6997 mL | 18.4986 mL | 36.9973 mL |
| | | 5 mM | 0.7399 mL | 3.6997 mL | 7.3995 mL |
| | 10 mM | 0.3700 mL | 1.8499 mL | 3.6997 mL | |
| <p>*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液: 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。</p> <p>储备液的保存方式和期限 -80°C, 6 months; -20°C, 1 month。 -80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。</p> | | | | | |
| References | <p>[1]. Garcia-Echeverria C, et al. Drug discovery approaches targeting the PI3K/Akt pathway in cancer. <i>Oncogene</i>. 2008 Sep 18;27(41):5511-26.</p> <p>[2]. Yu P, et al. Characterization of the activity of the PI3K/mTOR inhibitor XL765 (SAR245409) in tumor models with diverse genetic alterations affecting the PI3K pathway. <i>Mol Cancer Ther</i>. 2014 May;13(5):1078-91.</p> | | | | |
| | 实验参考: | | | | |
| Cell Assay | <p>Cellular proliferation is assessed using the Cell Proliferation ELISA, Bromodeoxyuridine Chemiluminescence Kit. Cytotoxicity is assessed using the ATP Bioluminescence Assay as follows: PC-3, MCF7, A549, LS174T, MDA-MB-468, U87-MG, and OVCAR-3 cells are plated at densities of 7×10^3, 1.5×10^4, 6×10^3, 7×10^3, 7×10^3, 6×10^3, 1.5×10^4 cells per well, respectively, onto 96-well microtiter plates in culture medium, incubated at 37°C, 5% CO₂ for 18 hours, and then treated with a serial dilution of compound in medium containing a final concentration of 0.3% DMSO. Triplicate wells are used for each compound concentration. Control wells receive 0.3% DMSO in media. Cultures are incubated at 37°C, 5% CO₂ for an additional 24 hours and cells are then assayed for viability using the ViaLight HS Kit[2].</p> | | | | |
| Animal Administration | <p>Mice[2]</p> <p>In vivo efficacy studies are performed in athymic nude mice. Tumor cells are cultured in DMEM supplemented with 10% FBS (20% for PC-3 and OVCAR-3 cells), Penicillin-Streptomycin, and nonessential amino acids at 37°C in a humidified 5% CO₂ atmosphere. On day 0, cells are harvested by brief trypsinization, and 1 to 5×10^6 cells in 0.1 mL ice-cold Hanks Balanced Salt Solution are implanted subcutaneously (OVCAR-3) or intradermally (MCF7 and U-87 MG) into the hind flank of female athymic nude mice. In the case of the MCF7 model, an estrogen pellet (IRA) is implanted subcutaneously at the nape of neck at the time of tumor cell implantation. A total of 3×10^6 PC-3 cells are similarly harvested and implanted subcutaneously into the hind-flank of 5- to 8-week-old male nude mice. Tumor growth is monitored weekly with calipers until staging and dose initiation. During the dosing period, body and tumor weights are assessed. Voxelalisib (XL-765) is formulated in sterile water/10 mM HCl or water and administered at the indicated doses and regimens by oral gavage at a dose volume of 10 mL/kg.</p> | | | | |
| References | <p>[1]. Garcia-Echeverria C, et al. Drug discovery approaches targeting the PI3K/Akt pathway in cancer. <i>Oncogene</i>. 2008 Sep 18;27(41):5511-26.</p> <p>[2]. Yu P, et al. Characterization of the activity of the PI3K/mTOR inhibitor XL765 (SAR245409) in tumor models with diverse genetic alterations affecting the PI3K pathway. <i>Mol Cancer Ther</i>. 2014</p> | | | | |



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

| |
|--------------------|
| May;13(5);1078-91. |
|--------------------|



源叶生物