



上海源叶生物科技有限公司
 Shanghai yuanye Bio-Technology Co., Ltd
 电话: 021-61312973 传真: 021-55068248
 网址: www.shyuanye.com
 邮箱: shyysw@sina.com


产品名称: **PAC-14028**
 产品别名: **Asivatrep**

| 生物活性: | | | | | |
|-------------------------------|---|-----------------------------------|-------------|-------------|--------------|
| Description | Asivatrep (PAC-14028) is a potent and selective transient receptor potential vanilloid type 1 (TRPV1) antagonist. | | | | |
| In Vitro | Asivatrep (PAC-14028) could prevent barrier damages, accelerate skin barrier recovery and suppress pruritus, showing a potential for the treatment of atopic dermatitis. It could suppress serum IgE increase, epidermal infiltration of inflammatory cells and mast cell degranulation associated with atopic dermatitis[1]. Asivatrep (PAC-14028) shows efficacies against diverse disease models including visceral pain, inflammatory bowel disease, and inflammatory pain[2]. | | | | |
| In Vivo | Asivatrep (PAC-14028) shows a plasma half-life of 2.1 h in rats while it is extended slightly to 3.8 h in minipigs. Oral bioavailability at 10 mg/kg dose is determined to be 52.7% and 64.2% in rats and minipigs, respectively suggesting that Asivatrep (PAC-14028) is relatively well-absorbed through oral route[1]. Asivatrep (PAC-14028) could inhibit capsaicin-evoked calcium influx in keratinocytes at sub-micromolar concentrations. This potent TRPV1 antagonistic activity in keratinocytes is manifested in vivo as the blockade of capsaicin-induced blood perfusion increase, and the accelerated barrier recovery from tape-stripping-induced barrier damages in hairless mice[3]. | | | | |
| Solvent&Solubility | In Vitro: DMSO : ≥ 28 mg/mL (56.97 mM) * "≥" means soluble, but saturation unknown. | | | | |
| | | Solvent Mass Concentration | 1 mg | 5 mg | 10 mg |
| | Preparing | 1 mM | 2.0347 mL | 10.1736 mL | 20.3471 mL |
| | Stock Solutions | 5 mM | 0.4069 mL | 2.0347 mL | 4.0694 mL |
| | | 10 mM | 0.2035 mL | 1.0174 mL | 2.0347 mL |
| | *请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液, 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限 -80°C, 6 months; -20°C, 1 month. -80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。 | | | | |
| References | [1]. Park YH, et al. Oral and topical pharmacokinetic studies of a novel TRPV1 antagonist, PAC-14028 in rats and minipigs using liquid chromatography/tandem mass spectrometric method. J Pharm Biomed Anal. 2012 Mar 5;61:8-14. [2]. Lim KM, et al. Development of PAC-14028, a novel transient receptor potential vanilloid type 1 (TRPV1) channel antagonist as a new drug for refractory skin diseases. Arch Pharm Res. 2012 Mar;35(3):393-6. [3]. Yun JW, et al. TRPV1 antagonist can suppress the atopic dermatitis-like symptoms by accelerating skin barrier recovery. J Dermatol Sci. 2011 Apr;62(1):8-15. | | | | |
| 实验参考: | | | | | |
| | Rats: 1 mg of Asivatrep (PAC-14028) is dissolved in 10 mL of methanol to obtain a concentration of 100 µg/mL. Male Sprague-Dawley rats/minipigs are given intravenously Asivatrep (PAC-14028) at a | | | | |



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

| | |
|------------------------------|--|
| Animal Administration | <p>single dose of 1 mg/kg, orally PAC-14028 at a single dose of 10 mg/kg as a suspension in 1% methylcellulose and 0.5% Tween80, or topically a single or multiple doses of 10 mg/kg as 1% Asivatrep (PAC-14028) solution (gel form) in 68% PEG400 vehicle. For the topical application, dorsal area is shaved and painted with designated dose of Asivatrep (PAC-14028) formulation. Occlusive dressing is placed on the applied region for 6 h to withhold licking or scratching the area. Blood samples are collected from the retroorbital sinus or jugular vein into heparinized tubes at designated times after drug administration. Blood samples are centrifuged immediately, and the plasma is collected and store for analysis[1].</p> |
| References | <p>[1]. Park YH, et al. Oral and topical pharmacokinetic studies of a novel TRPV1 antagonist, PAC-14028 in rats and minipigs using liquid chromatography/tandem mass spectrometric method. J Pharm Biomed Anal. 2012 Mar 5;61:8-14.</p> <p>[2]. Lim KM, et al. Development of PAC-14028, a novel transient receptor potential vanilloid type 1 (TRPV1) channel antagonist as a new drug for refractory skin diseases. Arch Pharm Res. 2012 Mar;35(3):393-6.</p> <p>[3]. Yun JW, et al. TRPV1 antagonist can suppress the atopic dermatitis-like symptoms by accelerating skin barrier recovery. J Dermatol Sci. 2011 Apr;62(1):8-15.</p> |



源叶生物