



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **BAY1125976**

产品别名: **BAY1125976**

生物活性:					
Description	BAY1125976 is a selective allosteric Akt1/Akt2 inhibitor; inhibits Akt1 and Akt2 activity with IC ₅₀ values of 5.2 nM and 18 nM at 10 μM ATP, respectively.				
IC ₅₀ & Target	Akt1	Akt2	Akt3		
	5.2 nM (IC ₅₀ , at 10 μM ATP)	18 nM (IC ₅₀ , at 10 μM ATP)	427 nM (IC ₅₀ , at 10 μM ATP)		
In Vitro	BAY 1125976 is equally potent against Akt1 (IC ₅₀ =5.2 nM at 10 μM ATP and 44 nM at 2 mM ATP) and Akt2 (IC ₅₀ =18 nM at 10 μM ATP and 36 nM at 2 mM ATP) isoforms and up to 86 fold less potent against Akt3 (IC ₅₀ =427 nM at 10 μM ATP). It inhibits the Akt1 and Akt2 by binding into an allosteric binding pocket formed by kinase and PH domain. It inhibits cell proliferation in a broad panel of human cancer cell lines, particularly in breast and prostate cancer cell lines expressing estrogen or androgen receptors. It effectively blocks Akt signaling by inhibiting the phosphorylation of Akt and the downstream effectors, including eukaryotic translation initiation factor 4E binding protein 1 (4E-BP1), glycogen synthase kinase 3 beta (GSK3s), proline-rich Akt substrate 40 kDa (PRAS40), S6 ribosomal protein (S6RP), and 70 kDa ribosomal protein S6 kinase 1 (70S6K) ^[1] .				
In Vivo	BAY 1125976 targets tumors displaying activation of the PI3K/Akt/mTOR pathway. BAY 1125976 exhibits strong <i>in vivo</i> efficacy in both cell line and patient-derived xenograft models such as the KPL4 breast cancer model (PIK3CA ^{H1074R} mutant), the MCF7 and HBCx-2 breast cancer models, and the Akt ^{E17K} mutant driven prostate cancer (LAPC-4) and anal cancer (AXF 984) models ^[1] .				
Solvent&Solubility	<i>In Vitro:</i> DMSO : 10.33 mg/mL (26.94 mM; Need ultrasonic and warming)				
		<div>Solvent / Mass / Concentration</div>	1 mg	5 mg	10 mg
	Preparing	1 mM	2.6079 mL	13.0395 mL	26.0790 mL
	Stock Solutions	5 mM	0.5216 mL	2.6079 mL	5.2158 mL
		10 mM	0.2608 mL	1.3040 mL	2.6079 mL
	<p>*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液; 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。</p> <p>储备液的保存方式和期限: -80℃, 6 months; -20℃, 1 month。 -80℃ 储存时, 请在 6 个月内使用, -20℃ 储存时, 请在 1 个月内使用。</p>				
References	[1]. Politz O, et al. BAY 1125976, a selective allosteric AKT1/2 inhibitor, exhibits high efficacy on AKT signaling-dependent tumor growth in mouse models. Int J Cancer. 2017 Jan 15;140(2):449-459.				
实验参考:					
Animal Administration	Mice: Female NMRI (nu/nu) mice s.c. injected with 3 x 10 ⁶ /100 μL KPL-4 breast cancer cells are used to study the mode-of-action of BAY 1125976. The treatment is started when tumors reaches 232-358 mm ³ in size and the mice receive a single oral dose of 25 or 50 mg/kg BAY 1125976. For determination of plasma concentration-time profiles, blood is drawn from the animals at different time points after compound administration ^[1] .				



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

Kinase Assay	PEG/water (60/40), pH 4.0, is used as a vehicle for BAY 1125976. The selectivity of BAY 1125976 is assessed using two different kinase panels: the 230 kinase panel; and the 468 kinase panel. In the 230 kinase panel, kinase activity is determined after incubation with 10 μ M BAY 1125976. An additional incubation with 1 μ M and 0.1 μ M BAY 1125976 is performed for the kinases where 10 μ M BAY 1125976 shows an inhibition over 70%. All tests are performed at 10 μ M ATP. The 468-kinase panel covered AGC, CAMK, CMGC, CK1, STE, TK, TKL, lipid, and atypical kinase families. The profiling is performed by combining the test compound with DNA-tagged kinase and immobilized ligands. The final kinase concentrations are measured by quantitative PCR ^[1] .
References	[1]. Politz O, et al. BAY 1125976, a selective allosteric AKT1/2 inhibitor, exhibits high efficacy on AKT signaling-dependent tumor growth in mouse models. Int J Cancer. 2017 Jan 15;140(2):449-459.

源叶生物