



上海源叶生物科技有限公司
 Shanghai yuanye Bio-Technology Co., Ltd
 电话: 021-61312973 传真: 021-55068248
 网址: www.shyuanye.com
 邮箱: shyysw@sina.com

产品名称: 葡萄糖酸锑钠
 产品别名: **Sodium stibogluconate; Stibogluconate trisodium nonahydrate**

生物活性:																											
Description	Sodium stibogluconate (Stibogluconate trisodium nonahydrate) is a potent inhibitor of protein tyrosine phosphatase. Sodium stibogluconate inhibits 99% of SHP-1, SHP-2 and PTP1B activity at 10, 100, 100 µg/mL, respectively.																										
IC₅₀ & Target	Phosphatase ^[1]																										
In Vitro	Sodium stibogluconate (Stibogluconate trisodium nonahydrate) inhibits 99% of SHP-1 activity at 10 µg/mL, a therapeutic concentration of the drug for leishmaniasis. Similar degrees of inhibition of SHP-2 and PTP1B required 100 µg/mL Sodium stibogluconate. The inhibition of cellular PTPases by the Sodium stibogluconate is suggested by its rapid induction of tyrosine phosphorylation of cellular proteins in Baf3 cells and its augmentation of IL-3-induced Janus family kinase 2/Stat5 tyrosine phosphorylation and proliferation of Baf3 cells. The augmentation of the opposite effects of GM-CSF and IFN-α on TF-1 cell growth by Sodium stibogluconate indicate its broad activities in the signaling of various cytokines ^[1] .																										
In Vivo	Sodium stibogluconate (Stibogluconate trisodium nonahydrate) induces 61% growth inhibition of Renca tumors in BALB/c mice coincident with an increase (2-fold) in tumor-infiltrating macrophages. A combination of Sodium stibogluconate and IL-2 is more effective in inhibiting tumor growth (91%) and inducing tumor-infiltrating (4-fold), whereas IL-2 alone has little effect ^[2] .																										
Solvent&Solubility	<p>In Vitro:</p> <p>H₂O : 9.1 mg/mL (9.99 mM; Need warming)</p> <p>H₂O : 6.67 mg/mL (7.32 mM; ultrasonic and adjust pH to 3 with 1M HCl)</p> <p>DMSO : < 1 mg/mL (insoluble or slightly soluble)</p>																										
	<table border="1"> <thead> <tr> <th rowspan="2"></th> <th colspan="2">Solvent</th> <th rowspan="2">1 mg</th> <th rowspan="2">5 mg</th> <th rowspan="2">10 mg</th> </tr> <tr> <th>Mass</th> <th>Concentration</th> </tr> </thead> <tbody> <tr> <td>Preparing</td> <td>1 mM</td> <td></td> <td>1.0978 mL</td> <td>5.4891 mL</td> <td>10.9782 mL</td> </tr> <tr> <td>Stock Solutions</td> <td>5 mM</td> <td></td> <td>0.2196 mL</td> <td>1.0978 mL</td> <td>2.1956 mL</td> </tr> <tr> <td></td> <td>10 mM</td> <td></td> <td>---</td> <td>---</td> <td>---</td> </tr> </tbody> </table>		Solvent		1 mg	5 mg	10 mg	Mass	Concentration	Preparing	1 mM		1.0978 mL	5.4891 mL	10.9782 mL	Stock Solutions	5 mM		0.2196 mL	1.0978 mL	2.1956 mL		10 mM		---	---	---
			Solvent					1 mg	5 mg	10 mg																	
		Mass	Concentration																								
Preparing	1 mM		1.0978 mL	5.4891 mL	10.9782 mL																						
Stock Solutions	5 mM		0.2196 mL	1.0978 mL	2.1956 mL																						
	10 mM		---	---	---																						
<p>*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液; 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。</p> <p>储备液的保存方式和期限 -80°C, 6 months; -20°C, 1 month。 -80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。</p>																											
References	<p>[1]. Pathak MK, et al. Sodium stibogluconate is a potent inhibitor of protein tyrosine phosphatases and augments cytokine responses in hemopoietic cell lines. J Immunol. 2001 Sep 15;167(6):3391-7.</p> <p>[2]. Fan K et al. Sodium Stibogluconate Interacts with IL-2 in Anti-Renca Tumor Action via a T Cell-Dependent Mechanism in Connection with Induction of Tumor-Infiltrating Macrophages. J Immunol. 2005 Nov 15;175(10):7003-8.</p>																										
实验参考:																											
	Human myeloid cell line TF-1 is maintained in RPMI 1640 supplemented with 10% FCS and 40																										



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

Cell Assay	ng/mL recombinant human GM-CSF. For cell proliferation assays, cells are washed in 10% FCS medium twice, resuspended in 10% FCS medium, incubated at 37°C for 16 h, and then cultured at 37°C in 10% FCS medium containing various amounts of cytokines, sodium stibogluconate, or potassium antimonyl tartrate for 3-6 days. The cell numbers in proliferation assays are determined by an MTT assay or by microscopic cell counting ^[1] .
Animal Administration	BALB/c and athymic nude BALB/c mice are inoculated (s.c.) at the flanks with Renca cells (106 cells/site). Four days after inoculation, the mice are subjected to no treatment (control) or treatment with IL-2 (105 IU/day for 5 days i.p.), Stibogluconate sodium (12 mg/day i.m. at hip regions), or the combination of the two agents for 2 wk. Tumor volume is measured during the study period and calculated using the formula for a prolate spheroid ^[2] .
References	[1]. Pathak MK, et al. Sodium stibogluconate is a potent inhibitor of protein tyrosine phosphatases and augments cytokine responses in hemopoietic cell lines. J Immunol. 2001 Sep 15;167(6):3391-7. [2]. Fan K et al. Sodium Stibogluconate Interacts with IL-2 in Anti-Renca Tumor Action via a T Cell-Dependent Mechanism in Connection with Induction of Tumor-Infiltrating Macrophages. J Immunol. 2005 Nov 15;175(10):7003-8.



源叶生物