



上海源叶生物科技有限公司  
 Shanghai yuanye Bio-Technology Co., Ltd  
 电话: 021-61312973 传真: 021-55068248  
 网址: www.shyuanye.com  
 邮箱: shyysw@sina.com

产品名称: **CCG215022**  
 产品别名: **CCG215022**

生物活性:					
<b>Description</b>	CCG215022 is a G protein-coupled receptor kinases (GRKs) inhibitor with IC <sub>50</sub> s of 0.15±0.07 μM, 0.38±0.06 μM and 3.9±1 μM for GRK2, GRK5 and GRK1, respectively.				
<b>IC<sub>50</sub> &amp; Target</b>	IC <sub>50</sub> & Target: 3.9±1.0 μM (GRK1), 0.15±0.07 μM (GRK2), 0.38±0.06 μM (GRK5), 120±40 μM (PKA) <sup>[1]</sup>				
<b>In Vitro</b>	CCG215022 has nanomolar potency against both GRK2 and GRK5 and is at least 20-fold more potent than Paroxetine. In the course of a GRK2 structure-based drug design campaign, CCG215022 exhibits nanomolar IC <sub>50</sub> values against both GRK2 and GRK5 and good selectivity against other closely related kinases such as GRK1 and PKA. Treatment of murine cardiomyocytes with CCG215022 results in significantly increases contractility at 20-fold lower concentrations than Paroxetine, an inhibitor with more modest selectivity for GRK2 <sup>[1]</sup> .				
<b>Solvent&amp;Solubility</b>	<b>In Vitro:</b> <b>DMSO : ≥ 28 mg/mL (56.06 mM)</b>  * "≥" means soluble, but saturation unknown.				
		<b>Solvent</b>	<b>Mass</b>		
		<b>Concentration</b>	<b>1 mg</b>	<b>5 mg</b>	<b>10 mg</b>
	<b>Preparing Stock Solutions</b>	1 mM	2.0020 mL	10.0100 mL	20.0200 mL
	5 mM	0.4004 mL	2.0020 mL	4.0040 mL	
	10 mM	0.2002 mL	1.0010 mL	2.0020 mL	
	*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液, 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80°C, 6 months; -20°C, 1 month。-80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。				
<b>References</b>	[1]. Homan KT, et al. Crystal Structure of G Protein-coupled Receptor Kinase 5 in Complex with a Rationally Designed Inhibitor. J Biol Chem. 2015 Aug 21;290(34):20649-59.				
实验参考:					
<b>Kinase Assay</b>	GRK5 and urea-washed bovine rod outer segments (ROS) are mixed in the dark in buffer containing 20 mM HEPES, pH 7.5, 4 mM MgCl <sub>2</sub> , and 2 mM EDTA and incubated for 35 min at room temperature. The reaction mixtures are exposed to ambient fluorescent light for 1 min prior to initiation of the reaction by addition of ATP (with [γ- <sup>32</sup> P]ATP) to a final concentration of 1 mM. Final concentration of GRK5 is 100 nM and ROS is between 0.75 and 24 μM. Reactions are initiated at room temperature, and samples are taken at 2-5 min and then quenched with SDS-PAGE loading dye. Proteins are separated using SDS-PAGE, gel is dried, and the incorporation of γ- <sup>32</sup> P is detected using a phosphor storage screen. Rates at 0 min are plotted against the ROS concentration, and V <sub>max</sub> and K <sub>m</sub> values are determined using the Michaelis-Menten equation. V <sub>max</sub> of each curve is normalized to the V <sub>max</sub> of GRK5561 run in parallel. Melting point determinations in response to 200 μM CCG215022 are performed in 20 mM HEPES, pH 7.0, 5 mM MgCl <sub>2</sub> , 2 mM DTT, 1 mM CHAPS at a final GRK5 concentration of 0.2 mg/mL and 100 μM anilinonaphthalene-8-sulfonic acid using a ThermoFluor plate reader. Melting points of GRK5 variants are				



上海源叶生物科技有限公司  
Shanghai yuanye Bio-Technology Co., Ltd  
电话: 021-61312973 传真: 021-55068248  
网址: [www.shyuanye.com](http://www.shyuanye.com)  
邮箱: [shyysw@sina.com](mailto:shyysw@sina.com)

	assayed in a buffer containing 20 mM HEPES, pH 8.0, 200 mM NaCl, 2 mM DTT, 2.5 mM MgCl <sub>2</sub> , and 0.1 mM anilinonaphthalene-8-sulfonic acid with or without 5 mM ATP. Final GRK5 concentration for these assays is 0.1 mg/mL <sup>[1]</sup> .
<b>References</b>	[1]. Homan KT, et al. Crystal Structure of G Protein-coupled Receptor Kinase 5 in Complex with a Rationally Designed Inhibitor. J Biol Chem. 2015 Aug 21;290(34):20649-59.



# 源叶生物