



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **BMT-145027**
产品别名: **BMT-145027**

生物活性:

Description	BMT-145027 is an mGluR5 positive allosteric modulator without inherent agonist activity, exhibits an EC50 of 47 nM.				
IC ₅₀ & Target	IC50: 47 nM (mGluR5)[1]				
In Vitro	BMT-145027 is a compound with high MsLM stability (85% remaining), acceptable potency (EC50=47 nM), and a modest decrease in planarity (Fsp3 = 0.17). Importantly, BMT-145027 lacks inherent mGluR5 agonist activity when tested at concentrations up to 16 μM[1].				
In Vivo	Drug-treated and control mice are shown two identical objects. After a 24-h natural forgetting period, the mice are reintroduced to a familiar object while simultaneously presented with a novel object. Since mice spend more time exploring unfamiliar objects, improved memory is measured as time spent exploring the novel object. BMT-145027 leads to a significant increase in time spent with the novel object when dosed at 30 mg/kg, with an apparent trend in novel object preference at 10 mg/kg. Satellite animals indicates a total plasma concentration of 2800 nM at 30 mg/kg[1].				
Solvent&Solubility	In Vitro: DMSO : 2.4 mg/mL (5.47 mM; Need ultrasonic and warming)				
	Preparing Stock Solutions	<div><div>Solvent</div><div>Mass</div><div>Concentration</div></div>	1 mg	5 mg	10 mg
		1 mM	2.2788 mL	11.3939 mL	22.7879 mL
		5 mM	0.4558 mL	2.2788 mL	4.5576 mL
		10 mM	---	---	---
	*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液。一旦配成溶液，请分装保存，避免反复冻融造成的产品失效。 储备液的保存方式和期限 -80℃, 6 months; -20℃, 1 month。 -80℃ 储存时，请在 6 个月内使用， -20℃ 储存时，请在 1 个月内使用。				
References	[1]. Hill MD, et al. Development of 1H-Pyrazolo[3,4-b]pyridines as Metabotropic Glutamate Receptor 5 Positive Allosteric Modulators. ACS Med Chem Lett. 2016 Oct 3;7(12):1082-1086.				

实验参考:

Animal Administration	Mice: Mice are dosed i.p. with either vehicle or BMT-145027 at 10 and 30 mg/kg 60 min prior to the training session and tested for recognition. Animal behavior is video recorded during both training and testing and the amount of time spent exploring the objects determined using Cleversys software[1].
Kinase Assay	Competition binding experiments are performed using a single concentration (5 nM) of [³ H]-MethoxyPyEP in the presence of increasing concentrations of BMT-145027. The reaction is terminated by the addition of 5 mL of ice-cold ish buffer and rapid filtration. The filter is punched into a 96 well flex-plate and scintillation cocktail is added to each well. The plate is allowed to soak for 8 h and then read on the micro-beta counter. IC ₅₀ values are determined using non-linear regression four-parameter logistic equation[1].



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

References

[1]. Hill MD, et al. Development of 1H-Pyrazolo[3,4-b]pyridines as Metabotropic Glutamate Receptor 5 Positive Allosteric Modulators. ACS Med Chem Lett. 2016 Oct 3;7(12):1082-1086.



源叶生物