



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: 他司鲁泰

产品别名: **Taspoglutide; ITM077; R1583; BIM51077**

生物活性:				
Description	Taspoglutide is a long-acting glucagon-like peptide 1 (GLP-1) receptor agonist developed for treatment of type 2 diabetes, with an EC50 value of 0.06 nM.			
IC₅₀ & Target	EC50: 0.06 nM (GLP-1)[1]			
In Vitro	Taspoglutide (R1583/BIM51077) is a long acting 10% formulation of (Aib8-35) human GLP-1 (7-36 amides) with 93% homology with the native polypeptide. It activates the GLP-1 receptor. Taspoglutide has comparable affinity (affinity constant 1.1±0.2 nM) to the natural ligand (affinity constant 1.5±0.3 nM) for the hGLP-1 receptor and exhibits comparable potency in stimulating cAMP production[2].			
In Vivo	Taspoglutide has been shown to enhance the rate of glucose-induced insulin secretion from isolated, cultured rat islets and the perfused ZDF rat pancreas. Taspoglutide in Sprague-Dawley rats and diabetic db/db mice have shown a dose-related enhancement of glucose-dependent insulin release, which lower blood glucose in the db/db mouse model of type 2 diabetes[3]. Acute treatment with taspoglutide reduces glucose excursion and increased insulin response during oGTT. In chronically treated rats, glucose excursion and levels of GIP, PYY and triglycerides during oGTT on day 21 are significantly reduced[4]. Hepatic triglyceride levels are significantly reduced in livers from taspoglutide-treated. Taspoglutide does not reduce plaque area or lipid content in the aortic arch or abdominal aorta, and no significant change in aortic macrophage accumulation is detected after taspoglutide or metformin mice[5].			
Solvent&Solubility	<i>In Vitro:</i> DMSO : ≥ 28 mg/mL (8.38 mM) * "≥" means soluble, but saturation unknown.			
		Solvent	Mass	
		Concentration		
	Preparing	1 mM	0.2994 mL	1.4971 mL
	Stock Solutions	5 mM	0.0599 mL	0.2994 mL
		10 mM	---	---
*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液; 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80°C, 6 months; -20°C, 1 month (protect from light)。-80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。				
References	<p>[1]. Sebkova E, et al. Taspoglutide, an analog of human glucagon-like Peptide-1 with enhanced stability and in vivo potency. Endocrinology. 2010 Jun;151(6):2474-82.</p> <p>[2]. Retterstol K, et al. Taspoglutide: a long acting human glucagon-like polypeptide-1 analogue. Expert Opin Investig Drugs. 2009 Sep;18(9):1405-11.</p> <p>[3]. Nauck MA, et al. Treatment with the human once-weekly glucagon-like peptide-1 analog taspoglutide in combination with metformin improves glycemic control and lowers body weight in patients with type 2 diabetes inadequately controlled with metformin alone: a double-blind placebo-controlled study. Diabetes</p>			



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

	<p>Care. 2009 Jul;32(7):1237-43.</p> <p>[4]. Sebkova E, et al. Taspoglutide, a novel human once-weekly analogue of glucagon-like peptide-1, improves glucose homeostasis and body weight in the Zucker diabetic fatty rat. Diabetes Obes Metab. 2010 Aug;12(8):674-82.</p> <p>[5]. Panjwani N, et al. GLP-1 receptor activation indirectly reduces hepatic lipid accumulation but does not attenuate development of atherosclerosis in diabetic male ApoE(-/-) mice. Endocrinology. 2013 Jan;154(1):127-39.</p>
实验参考:	
Animal Administration	<p>Rats: The Zucker diabetic fatty (ZDF) rats (animal model of type 2 diabetes) are used in the study. ZDF rats are treated acutely (0.1, 1, 10 µg/kg) or chronically (sustained-release of 1 mg) with a single long-acting dose of taspoglutide. Pioglitazone is used as a positive control in the chronic study. Postprandial glucose, body weight, glycaemic control and insulin sensitivity are assessed over 21 days in chronically treated animals[4].</p> <p>Mice: High-fat diet-fed male ApoE^{-/-} mice are used in the study. Mice with glucose levels from 15-25 mM are then randomized to different groups and treated for 12 wk with a once-monthly sc 0.4-mg taspoglutide microtablet suspension, a sc placebo microtablet, or metformin (400 mg/kg*d) continuously provided in the drinking water plus a sc placebo microtablet[5].</p>
References	<p>[1]. Sebkova E, et al. Taspoglutide, an analog of human glucagon-like Peptide-1 with enhanced stability and in vivo potency. Endocrinology. 2010 Jun;151(6):2474-82.</p> <p>[2]. Retterstol K, et al. Taspoglutide: a long acting human glucagon-like polypeptide-1 analogue. Expert Opin Investig Drugs. 2009 Sep;18(9):1405-11.</p> <p>[3]. Nauck MA, et al. Treatment with the human once-weekly glucagon-like peptide-1 analog taspoglutide in combination with metformin improves glycemic control and lowers body weight in patients with type 2 diabetes inadequately controlled with metformin alone: a double-blind placebo-controlled study. Diabetes Care. 2009 Jul;32(7):1237-43.</p> <p>[4]. Sebkova E, et al. Taspoglutide, a novel human once-weekly analogue of glucagon-like peptide-1, improves glucose homeostasis and body weight in the Zucker diabetic fatty rat. Diabetes Obes Metab. 2010 Aug;12(8):674-82.</p> <p>[5]. Panjwani N, et al. GLP-1 receptor activation indirectly reduces hepatic lipid accumulation but does not attenuate development of atherosclerosis in diabetic male ApoE(-/-) mice. Endocrinology. 2013 Jan;154(1):127-39.</p>