



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: 角黄素

产品别名: **Canthaxanthin; E 161g; all-trans-Canthaxanthin**

| | | | | |
|--|--|-------------------------------------|-----------|-----------|
| 生物活性: | | | | |
| Description | Canthaxanthin is a red-orange carotenoid with various biological activities, such as antioxidant, antitumor properties. | | | |
| In Vitro | Canthaxanthin enrichment of LDL has the potential to protect cholesterol from oxidation. In addition to its free radical scavenging and antioxidant properties (e.g., the induction of catalase and superoxide dismutase), canthaxanthin shows immunomodulatory activity (e.g. enhancing the proliferation and function of immune competent cells) and plays important role in gap junction communication (e.g. induction of the transmembrane protein connexin)[1]. At concentrations of 0.1 to 1 x 1000 µM, canthaxanthin significantly reduces the overall number of tumor cells. The greatest inhibition is observed at a canthaxanthin concentration of 1000 after 72 h and 96 h of incubation[2]. | | | |
| In Vivo | Canthaxanthin alters the protective ability of tissues against oxidative stress. Canthaxanthin treatment for 15 d at the dose of 14 µg/kg body weight results in hepatic incorporation of the carotenoid, which is maximum in liver and reaches 0.52 ± 0.05 nmol/g liver. Glutathione peroxidase activity is 35% lower and catalase (59%) and manganese superoxide dismutase (28%) activities are higher in canthaxanthin-treated mice than in controls[3]. Canthaxanthin inhibit the growth of mammary tumors in mice and the anti-tumor activity is also influenced by the supplemental dose[4]. Diet supplementation with canthaxanthin for 3 weeks prior to the carcinogen results in a 65% reduction in the number of mammary cancers by a mechanism not involving pro-vitamin A activity[5]. | | | |
| Solvent&Solubility | In Vitro: DMSO : 5 mg/mL (8.85 mM; Need ultrasonic and warming) | | | |
| | | <div>SolventMassConcentration</div> | 1 mg | 5 mg |
| | Preparing | 1 mM | 1.7704 mL | 8.8521 mL |
| | Stock Solutions | 5 mM | 0.3541 mL | 1.7704 mL |
| | | 10 mM | --- | --- |
| *请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液。一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80°C, 6 months; -20°C, 1 month (protect from light, stored under nitrogen)。-80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。 | | | | |
| References | [1]. Esatbeyoglu T, et al. Canthaxanthin: From molecule to function. Mol Nutr Food Res. 2017 Jun;61(6). [2]. Huang DS, et al. Inhibitory effects of canthaxanthin on in vitro growth of murine tumor cells. Cancer Lett. 1992 Aug 31;65(3):209-13. [3]. Palozza P, et al. Canthaxanthin supplementation alters antioxidant enzymes and iron concentration in liver of Balb/c mice. J Nutr. 2000 May;130(5):1303-8. [4]. Chew BP, et al. A comparison of the anticancer activities of dietary beta-carotene, canthaxanthin and astaxanthin in mice in vivo. Anticancer Res. 1999 May-Jun;19(3A):1849-53. | | | |



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

| | |
|------------------------------|--|
| | [5]. Grubbs CJ, et al. Effect of canthaxanthin on chemically induced mammary carcinogenesis. Oncology. 1991;48(3):239-45. |
| 实验参考: | |
| Animal Administration | Mice: Female 6-wk-old Balb/c mice are randomly divided into two groups (n = 10/group). The control group receives olive oil alone (vehicle) and the canthaxanthin-treated group receives canthaxanthin at a dose of 14 µg/kg body weight per day. At the end of the treatment, mice are killed by cervical dislocation and liver is excised, frozen in liquid nitrogen and stored at -80°C[3]. |
| References | <p>[1]. Esatbeyoglu T, et al. Canthaxanthin: From molecule to function. Mol Nutr Food Res. 2017 Jun;61(6).</p> <p>[2]. Huang DS, et al. Inhibitory effects of canthaxanthin on in vitro growth of murine tumor cells. Cancer Lett. 1992 Aug 31;65(3):209-13.</p> <p>[3]. Palozza P, et al. Canthaxanthin supplementation alters antioxidant enzymes and iron concentration in liver of Balb/c mice. J Nutr. 2000 May;130(5):1303-8.</p> <p>[4]. Chew BP, et al. A comparison of the anticancer activities of dietary beta-carotene, canthaxanthin and astaxanthin in mice in vivo. Anticancer Res. 1999 May-Jun;19(3A):1849-53.</p> <p>[5]. Grubbs CJ, et al. Effect of canthaxanthin on chemically induced mammary carcinogenesis. Oncology. 1991;48(3):239-45.</p> |

源叶生物