



上海源叶生物科技有限公司
 Shanghai yuanye Bio-Technology Co., Ltd
 电话: 021-61312973 传真: 021-55068248
 网址: www.shyuanye.com
 邮箱: shyysw@sina.com

产品名称: **SJB2-043**
 产品别名: **SJB2-043**

生物活性:					
Description	SJB2-043 is an inhibitor of the native USP1/UAF1 complex with IC ₅₀ of 544 nM.				
IC₅₀ & Target	IC ₅₀ : 544 nM (USP1/UAF1)[1]				
In Vitro	SJB2-043 causes a dose-dependent decrease in ubiquitin-specific protease 1 (USP1) levels and a concomitant degradation of inhibitor of DNA-binding-1 (ID1) protein in the K562 cells at a micromolar drug concentration. SJB2-043 also causes a decrease in the levels of other ID proteins, namely ID2 and ID3 in K562 cells. SJB2-043 causes a dose-dependent decrease in the number of viable K562 cells, with an EC ₅₀ of approximately 1.07 μM. Moreover, SJB2-043 induces apoptosis of K562 cells in a dose-dependent manner[1].				
Solvent&Solubility	In Vitro: DMSO : 3.33 mg/mL (12.10 mM; Need ultrasonic) H ₂ O : < 0.1 mg/mL (insoluble)				
		Solvent Mass Concentration	1 mg	5 mg	10 mg
	Preparing	1 mM	3.6329 mL	18.1646 mL	36.3293 mL
	Stock Solutions	5 mM	0.7266 mL	3.6329 mL	7.2659 mL
		10 mM	0.3633 mL	1.8165 mL	3.6329 mL
	*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液。一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80°C, 6 months; -20°C, 1 month。-80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。				
References	[1]. Mistry H, et al. Small molecule inhibitors of USP1 target ID1 degradation in leukemic cells. Mol Cancer Ther. 2013 Dec;12(12):2651-62.				
实验参考:					
Cell Assay	Leukemic cell lines are grown in RPMI 1640 medium supplemented with 10% fetal bovine serum and penicillin/streptomycin. Hela cells and U2OS cells are grown in DMEM supplemented with 10% fetal bovine serum and penicillin/streptomycin. USP1 inhibitor C527 and its derivatives (e.g., SJB2-043) are synthesized and the purity is validated by high-performance liquid chromatography. Primary human AML patient samples are collected from DFCI leukemia program under the approval of appropriate protocols. Cells are treated with DMSO or USP1 inhibitors (e.g., SJB2-043) in appropriate medium for 24-72 hrs. The viable cell counts are determined using Trypan blue staining, Cell TiterGlo reagent or MTT assay. The apoptotic cells are detected using AnnexinV and 7AAD staining using flow cytometry. For Benzidine staining, the cells are washed twice with PBS and resuspended in 45 μL of PBS + 5 μL of Benzidine stain solution (0.2% in 0.5 M glacial acetic acid, 3% H ₂ O ₂). After 45 min incubation at room temperature, the Benzidine positive cells are detected by light microscopy[1].				
	The in vitro enzymatic assays are performed using ubiquitin-AMC (Ub-7-amido-4methylcoumarin) as a substrate in a reaction buffer containing 20 mM HEPES-KOH (pH 7.8), 20 mM NaCl, 0.1 mg/mL ovalbumin,				



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

Kinase Assay	0.5 mM EDTA and 10 mM dithiothreitol. The fluorescence is measured by FluoStar Galaxy Fluorometer. For the Ub-vinylsulfone (VS) assay, the proteins are incubated with Ub-VS at 0.5 μ M final concentration for 45 min at 30°C, followed by the immunoblotting analysis[1].
References	[1]. Mistry H, et al. Small molecule inhibitors of USP1 target ID1 degradation in leukemic cells. Mol Cancer Ther. 2013 Dec;12(12):2651-62.



源叶生物