



上海源叶生物科技有限公司
 Shanghai yuanye Bio-Technology Co., Ltd
 电话: 021-61312973 传真: 021-55068248
 网址: www.shyuanye.com
 邮箱: shyysw@sina.com

产品名称: **Reversine**
 产品别名: 逆转素

生物活性:					
Description	Reversine is a novel class of ATP-competitive Aurora kinase inhibitor with IC ₅₀ s of 400, 500 and 400 nM for Aurora A, Aurora B and Aurora C, respectively.				
IC₅₀ & Target	Aurora A	Aurora B	Aurora C		
	400 nM (IC ₅₀)	500 nM (IC ₅₀)	400 nM (IC ₅₀)		
In Vitro	Reversine, a novel Aurora kinases inhibitor, inhibits colony formation of human acute myeloid leukemia cells. Reversine is a potent inhibitor of Aurora A and B and is also an inhibitor of Aurora C kinase. Aurora A and B activities are inhibited by 80% and Aurora kinase C by 55%, already at a concentration of 0.5 μM, whereas no inhibition or only modest inhibition is observed on others kinases tested. In a second round of experiments, the IC ₅₀ of Reversine is determined on Aurora kinase A to be 400 nM, whereas Aurora kinase B and C IC ₅₀ are 500 and 400 nM, respectively. The IC ₅₀ is also determined on MEK1 is >1.5 μM and that the IC ₅₀ on muscle myosin (an analogue of nonmuscle myosin II) is 350 nM[1].				
In Vivo	The combination of Reversine and aspirin can more efficiently induce cell cycle arrest and apoptosis. To evaluate the anti-tumor effect of this combination, a xenograft nude mouse model is established by s.c. injection. Mice inoculated with cervical cancer cells have lost about 10 % of their initial body weight by about 16 days after tumor inoculation. However, tumor growth (tumor weight) is reduced and the mice survive longer in the combination group[2].				
Solvent&Solubility	In Vitro: DMSO : 20 mg/mL (50.83 mM; Need ultrasonic) H ₂ O : < 0.1 mg/mL (insoluble)				
		Solvent Mass Concentration	1 mg	5 mg	10 mg
	Preparing	1 mM	2.5414 mL	12.7068 mL	25.4136 mL
	Stock Solutions	5 mM	0.5083 mL	2.5414 mL	5.0827 mL
		10 mM	0.2541 mL	1.2707 mL	2.5414 mL
*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液,一旦配成溶液,请分装保存,避免反复冻融造成的产品失效。 储备液的保存方式和期限 -80°C, 6 months; -20°C, 1 month。-80°C 储存时,请在 6 个月内使用, -20°C 储存时,请在 1 个月内使用。					
References	[1]. D'Alise AM, et al. Reversine, a novel Aurora kinases inhibitor, inhibits colony formation of human acute myeloid leukemia cells. Mol Cancer Ther. 2008 May;7(5):1140-9. [2]. Qin HX, et al. Synergistic antitumor activity of reversine combined with aspirin in cervical carcinoma in vitro and in vivo. Cytotechnology. 2013 Aug;65(4):643-53.				
实验参考:					
	HCT116 and HL60 cells are incubated with either 5 μmol/L Reversine or DMSO 0.01%. Cells are harvested and fixed in 70% ethanol overnight. After double washing with PBS, cells are labeled with cell cycle staining reagent PBS, 0.1% Triton X-100, 200 μg/mL DNase-free RNase, and 25 μg/mL				



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

Cell Assay	<p>propidium iodide and incubated at room temperature in the dark for 30 min. DNA content is analyzed using FACSCalibur. Cell viability of different tumor cell lines is assessed using ATPlite 1step. Briefly, 2×10^4 cells for each well are plated in a 96-well plate in presence of crescent quantity of Reversine. After 72 h, the plates are recovered and 100 μL ATPlite solution is added to each well. The plates are shaken for 2 min at 700 rpm and luminescence is measured using EnVision Multilabel plate reader. Each sample is analyzed in triplicate[1].</p>
Animal Administration	<p>Mice[2] Female athymic 6-8 weeks old BALB/c nude mice are used. U14 cell suspension (5×10^6 cells in 100 μL of RPMI-1640 medium) is injected subcutaneously. The purpose of developing cervical tumors is to generate histological intact tumors for drug therapy. When the diameter of tumors reached up to about 1 cm, Reversine, aspirin or their combinations are administrated by intraperitoneal injection per 3 days, twenty-five of these mice are randomly assigned into one of the following five groups: (a) mice treated with RPMI-1640 medium, (b) mice treated with DMSO, (c) mice treated with Reversine (10 mg/kg), (d) mice treated with aspirin (1 μg/kg) and (e) mice treated with a Reversine and aspirin combination. Body weight and tumor size at the site of inoculation are measured three times a week. Tumor size is measured every 3 days from two diameters, tumor volume is estimated using the formula $L \times S^2 / 2$ (L as the longest diameter, S as the shortest diameter).</p>
References	<p>[1]. D'Alise AM, et al. Reversine, a novel Aurora kinases inhibitor, inhibits colony formation of human acute myeloid leukemia cells. Mol Cancer Ther. 2008 May;7(5):1140-9. [2]. Qin HX, et al. Synergistic antitumor activity of reversine combined with aspirin in cervical carcinoma in vitro and in vivo. Cytotechnology. 2013 Aug;65(4):643-53.</p>

源叶生物