



上海源叶生物科技有限公司
 Shanghai yuanye Bio-Technology Co., Ltd
 电话: 021-61312973 传真: 021-55068248
 网址: www.shyuanye.com
 邮箱: shyysw@sina.com

产品名称: **Bitopertin**
 产品别名: **RG1678; RO4917838**

生物活性:																														
Description	Bitopertin is a potent, noncompetitive glycine reuptake inhibitor, inhibits glycine uptake at human GlyT1 with a concentration exhibiting IC50 of 25 nM.																													
IC₅₀ & Target	IC50: 25 nM (GlyT1)[1]																													
In Vitro	Bitopertin (RG1678) competitively blocks [³ H]ORG24598 binding sites at human GlyT1b in membranes from Chinese hamster ovary cells. Bitopertin potently inhibits [³ H]glycine uptake in cells stably expressing hGlyT1b and mGlyT1b, with IC ₅₀ values of 25±2 nM and 22±5 nM, respectively (n=6). Conversely, Bitopertin has no effect on hGlyT2-mediated glycine uptake up to 30 μM concentration. Bitopertin has high affinity for the recombinant hGlyT1b transporter. Under equilibrium conditions (1 h at room temperature), Bitopertin displaces [³ H]ORG24598 binding with a K _i of 8.1 nM. In hippocampal CA1 pyramidal cells, Bitopertin enhances NMDA-dependent long-term potentiation at 100 nM but not at 300 nM ^[1] . Additional profiling revealed that Bitopertin (RG1678) has an excellent selectivity profile against the GlyT2 isoform (IC ₅₀ >30 μM) and toward a panel of 86 targets including transmembrane and soluble receptors, enzymes, ion channels, and monoamine transporters (<41% inhibition at 10 μM is measured for all targets)[2].																													
In Vivo	Bitopertin (RG1678) dose-dependently increases cerebrospinal fluid and striatal levels of glycine measured by microdialysis in rats. Additionally Bitopertin attenuates hyperlocomotion induced by the psychostimulant D-amphetamine or the NMDA receptor glycine site antagonist L-687,414 in mice. Bitopertin also prevents the hyper-response to D-amphetamine challenge in rats treated chronically with phencyclidine, an NMDA receptor open-channel blocker. Administration of vehicle has no effect on extracellular levels of striatal glycine, which remained constant throughout the experiment. In contrast, p.o. administration of Bitopertin (1-30 mg/kg) produced a dose-dependent increase in extracellular glycine levels. Bitopertin 30 mg/kg produces glycine levels 2.5 times higher than pretreatment levels. A similar dose-dependent increase in glycine concentration is observed in the CSF of rats treated p.o. with Bitopertin (1-10 mg/kg) compared with vehicle-treated animals, 3 h after drug administration. Interestingly, the level of CSF glycine increase 3 h after Bitopertin dosing is very similar to the increase in the microdialysis experiment at the same time point[1]. In vivo pharmacokinetic studies in rat and monkey reveals that Bitopertin (RG1678) has, in both species, a low plasma clearance, an intermediate volume of distribution, a good oral bioavailability (78% for rat, 56% for monkey), and a favorable terminal half-life (5.8 h for rat, 6.4 h for monkey). The plasma protein binding is high in the two preclinical species (97%) and in human (98%). The CNS penetration of Bitopertin in rat (brain/plasma=0.7) is better than that in mouse (brain/plasma=0.5)[2].																													
	<p><i>In Vitro:</i></p> <p>DMSO : ≥ 50 mg/mL (92.00 mM)</p> <p>* "≥" means soluble, but saturation unknown.</p> <table border="1"> <thead> <tr> <th rowspan="2">Preparing Stock Solutions</th> <th>Solvent</th> <th>Mass</th> <th rowspan="2">1 mg</th> <th rowspan="2">5 mg</th> <th rowspan="2">10 mg</th> </tr> <tr> <th colspan="2">Concentration</th> </tr> </thead> <tbody> <tr> <td></td> <td>1 mM</td> <td></td> <td>1.8401 mL</td> <td>9.2003 mL</td> <td>18.4006 mL</td> </tr> <tr> <td></td> <td>5 mM</td> <td></td> <td>0.3680 mL</td> <td>1.8401 mL</td> <td>3.6801 mL</td> </tr> <tr> <td></td> <td>10 mM</td> <td></td> <td>0.1840 mL</td> <td>0.9200 mL</td> <td>1.8401 mL</td> </tr> </tbody> </table>				Preparing Stock Solutions	Solvent	Mass	1 mg	5 mg	10 mg	Concentration			1 mM		1.8401 mL	9.2003 mL	18.4006 mL		5 mM		0.3680 mL	1.8401 mL	3.6801 mL		10 mM		0.1840 mL	0.9200 mL	1.8401 mL
Preparing Stock Solutions	Solvent	Mass	1 mg	5 mg		10 mg																								
	Concentration																													
	1 mM		1.8401 mL	9.2003 mL	18.4006 mL																									
	5 mM		0.3680 mL	1.8401 mL	3.6801 mL																									
	10 mM		0.1840 mL	0.9200 mL	1.8401 mL																									



上海源叶生物科技有限公司
 Shanghai yuanye Bio-Technology Co., Ltd
 电话: 021-61312973 传真: 021-55068248
 网址: www.shyuanye.com
 邮箱: shyysw@sina.com

<p>Solvent&Solubility</p>	<p>*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液; 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。</p> <p>储备液的保存方式和期限: -80°C, 6 months; -20°C, 1 month。-80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。</p> <p>In Vivo:</p> <p>请根据您的实验动物和给药方式选择适当的溶解方案。以下溶解方案都请先按照 In Vitro 方式配制澄清的储备液, 再依次添加助溶剂:</p> <p>——为保证实验结果的可靠性, 澄清的储备液可以根据储存条件, 适当保存: 体内实验的工作液, 建议您现用现配, 当天使用; 以下溶剂前显示的百分比是指该溶剂在您配制终溶液中的体积占比; 如在配制过程中出现沉淀、析出现象, 可以通过加热和/或超声的方式助溶</p> <p>1.请依序添加每种溶剂: 10% DMSO→40% PEG300 →5% Tween-80 → 45% saline</p> <p>Solubility: ≥ 2.5 mg/mL (4.60 mM); Clear solution</p> <p>此方案可获得 ≥ 2.5 mg/mL (4.60 mM, 饱和度未知) 的澄清溶液。</p> <p>以 1 mL 工作液为例, 取 100 μL 25.0 mg/mL 的澄清 DMSO 储备液加到 400 μL PEG300 中, 混合均匀向上述体系中加入 50 μL Tween-80, 混合均匀; 然后继续加入 450 μL 生理盐水定容至 1 mL。</p> <p>2.请依序添加每种溶剂: 10% DMSO→ 90% (20% SBE-β-CD in saline)</p> <p>Solubility: ≥ 2.5 mg/mL (4.60 mM); Clear solution</p> <p>此方案可获得 ≥ 2.5 mg/mL (4.60 mM, 饱和度未知) 的澄清溶液。</p> <p>以 1 mL 工作液为例, 取 100 μL 25.0 mg/mL 的澄清 DMSO 储备液加到 900 μL 20% 的 SBE-β-CD 生理盐水溶液中, 混合均匀。</p> <p>3.请依序添加每种溶剂: 10% DMSO →90% corn oil</p> <p>Solubility: ≥ 2.5 mg/mL (4.60 mM); Clear solution</p> <p>此方案可获得 ≥ 2.5 mg/mL (4.60 mM, 饱和度未知) 的澄清溶液, 此方案不适用于实验周期在半个月以上的实验。</p> <p>以 1 mL 工作液为例, 取 100 μL 25.0 mg/mL 的澄清 DMSO 储备液加到 900 μL 玉米油中, 混合均匀。</p>
<p>References</p>	<p>[1]. Alberati D, et al. Glycine reuptake inhibitor RG1678: A pharmacologic characterization of an investigational agent for the treatment of schizophrenia. <i>Neuropharmacology</i>. 2012 Feb;62(2):1152-61.</p> <p>[2]. Pinard E, et al. Selective GlyT1 Inhibitors: Discovery of [4-(3-Fluoro-5-trifluoromethylpyridin-2-yl)piperazin-1-yl][5-methanesulfonyl-2-((S)-2,2,2-trifluoro-1-methyl ethoxy)phenyl]methanone (RG1678), a Promising Novel Medicine To Treat Schizophrenia. <i>J Me</i></p>
<p>实验参考:</p>	
	<p>Mice[1]</p> <p>Male NMRI mice (20-30 g) are treated with Bitopertin (0.3, 3, 1, and 10 mg/kg p.o.) or vehicle (p.o.). After 1 min, L-687,414 (50 mg/kg s.c.) or vehicle is given. After 15 min of habituation in the activity chambers, horizontal activity is recorded for 60 min. The time course of Bitopertin effects on L-678,414-induced hyperactivity is also examined; locomotor activity is assessed 2.5, 4.5, and 24 h after administration of Bitopertin (L-678,414 is always given 15 min before the activity procedure). In addition, the effect of subchronic Bitopertin is investigated. Mice receive vehicle or Bitopertin (1</p>



上海源叶生物科技有限公司
 Shanghai yuanye Bio-Technology Co., Ltd
 电话: 021-61312973 传真: 021-55068248
 网址: www.shyuanye.com
 邮箱: shyysw@sina.com

<p>Animal Administration</p>	<p>mg/kg p.o.) for 4 consecutive days and L-678,414-induced hyperactivity is evaluated on day 5.</p> <p>Rats[1]</p> <p>Wistar rats receive a 14-day treatment of PCP HC1 (5 mg/kg) or vehicle (NaCl 0.9%, 5 mL/kg i.p.). 24 h following the last injection, rats (6-18 per group) are allowed to individually habituate to the test boxes for 30 min. Rats then received Bitopertin (1, 3, 10 mg/kg p.o.) or vehicle (Polysorbate 80, HEC, Methyl- and Propylparaben pH 6.0; 5 mL/kg p.o.), followed after 1 h by 1 mg/kg D-amphetamine or vehicle i.p. Horizontal activity is recorded directly after the administration of Bitopertin until 120 min after dosing with amphetamine. Data are analyzed by ANOVA supplemented by Fischer's least significant difference post hoc test.</p>
<p>Kinase Assay</p>	<p>Association and dissociation kinetic analysis of [³H]ORG24598 to hGlyT1 and ratforebrain membranes is performed. [³H]ORG24598 binding experiments are performed using membranes from CHO cells expressing hGlyT1b and also in membranes from mouse, rat, monkey, and dogforebrains. Saturation isotherms are determined by adding [³H]ORG24598 to rat, mouse, monkey, and dog forebrain membranes (40 µg/well) and cell membranes (10 µg/well) in a total volume of 500 µL for 3 h at room temperature. Saturation binding experiments are analyzed by an Excel-based curve-fitting program using the Michaelis-Menten equation derived from the equation of a bimolecular reaction and the law of mass action: $B = (B_{max} \times [F]) / (K_d + [F])$, where B is the amount of ligand bound at equilibrium, B_{max} the maximum number of binding sites, [F] the concentration of free ligand, and K_d the ligand dissociation constant. For inhibition experiments, membranes are incubated with 3 nM [³H]ORG24598 and 10 concentrations of Bitopertin for 1 h at room temperature. Schild analysis is performed in the presence of increasing concentrations of [³H]ORG24598 (1-300 nM). IC_{50} values are derived as described above. K_i values are calculated according to the following equation: $K_i = IC_{50} / (1 + [L] / K_d)$ [1].</p>
<p>References</p>	<p>[1]. Alberati D, et al. Glycine reuptake inhibitor RG1678: A pharmacologic characterization of an investigational agent for the treatment of schizophrenia. <i>Neuropharmacology</i>. 2012 Feb;62(2):1152-61.</p> <p>[2]. Pinard E, et al. Selective GlyT1 Inhibitors: Discovery of [4-(3-Fluoro-5-trifluoromethylpyridin-2-yl)piperazin-1-yl][5-methanesulfonyl-2-((S)-2,2,2-trifluoro-1-methylethoxy)phenyl]methanone (RG1678), a Promising Novel Medicine To Treat Schizophrenia. <i>J Me</i></p>