



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **Setiptiline (maleate)**

产品别名: 马来酸司普替林; **MO-8282**

生物活性:				
Description	Setiptiline is a serotonin receptor antagonist. IC50 value: Target: 5-HT receptor Setiptiline is a tetracyclic antidepressant (TeCA) which acts as a noradrenergic and specific serotonergic antidepressant (NaSSA). Setiptiline acts as a norepinephrine reuptake inhibitor, α 2-adrenergic receptor antagonist, and serotonin receptor antagonist, likely at the 5-HT2A, 5-HT2C, and/or 5-HT3 subtypes, as well as an H1 receptor inverse agonist/antihistamine. From Wikipedia.			
Solvent&Solubility	In Vitro: DMSO : 100 mg/mL (264.95 mM; Need ultrasonic)			
	Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg
		1 mM	2.6495 mL	13.2475 mL
		5 mM	0.5299 mL	2.6495 mL
		10 mM	0.2649 mL	1.3247 mL
	*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液; 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80°C, 6 months; -20°C, 1 month. -80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。			
	In Vivo: 请根据您的实验动物和给药方式选择适当的溶解方案。以下溶解方案都请先按照 In Vitro 方式配制澄清的储备液, 再依次添加助溶剂: ——为保证实验结果的可靠性, 澄清的储备液可以根据储存条件, 适当保存; 体内实验的工作液, 建议您现用现配, 当天使用; 以下溶剂前显示的百分比是指该溶剂在您配制终溶液中的体积占比; 如在配制过程中出现沉淀、析出现象, 可以通过加热和/或超声的方式助溶 1.请依序添加每种溶剂: 10% DMSO→40% PEG300 →5% Tween-80 → 45% saline Solubility: ≥ 2.5 mg/mL (6.62 mM); Clear solution 此方案可获得 ≥ 2.5 mg/mL (6.62 mM, 饱和度未知) 的澄清溶液。 以 1 mL 工作液为例, 取 100 μ L 25.0 mg/mL 的澄清 DMSO 储备液加到 400 μ L PEG300 中, 混合均匀, 向上述体系中加入 50 μ L Tween-80, 混合均匀; 然后继续加入 450 μ L 生理盐水定容至 1 mL。 2.请依序添加每种溶剂: 10% DMSO→ 90% (20% SBE- β -CD in saline) Solubility: ≥ 2.5 mg/mL (6.62 mM); Clear solution 此方案可获得 ≥ 2.5 mg/mL (6.62 mM, 饱和度未知) 的澄清溶液。 以 1 mL 工作液为例, 取 100 μ L 25.0 mg/mL 的澄清 DMSO 储备液加到 900 μ L 20% 的 SBE- β -CD 生理盐水水溶液中, 混合均匀。 3.请依序添加每种溶剂: 10% DMSO →90% corn oil Solubility: ≥ 2.5 mg/mL (6.62 mM); Clear solution 此方案可获得 ≥ 2.5 mg/mL (6.62 mM, 饱和度未知) 的澄清溶液, 此方案不适用于实验周期在半个月以上的			



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

	<p>实验。</p> <p>以 1 mL 工作液为例, 取 100 μL 25.0 mg/mL 的澄清 DMSO 储备液加到 900 μL 玉米油中, 混合均匀。</p>
References	<p>[1]. Niho T, et al. [Pharmacological properties of MO-8282, a novel antidepressant]. Nihon Yakurigaku Zasshi. 1986 Oct;88(4):309-20.</p> <p>[2]. Setiptiline</p> <p>[3]. Yamada K, et al. [Behavioral effects of a new antidepressant, setiptiline]. Nihon Yakurigaku Zasshi. 1991 Jan;97(1):31-9.</p> <p>[4]. Kamimura M, et al. The effect of age on plasma level of setiptiline maleate in depressed patients. Prog Neuropsychopharmacol Biol Psychiatry. 1994 Oct;18(6):1015-26.</p> <p>[5]. Kuniyoshi M, et al. Effectiveness of concomitant setiptiline maleate (Tecipul) on negative symptoms of schizophrenia. Prog Neuropsychopharmacol Biol Psychiatry. 1994 Mar;18(2):339-46.</p>

源叶生物